

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number  
**WO 02/24881 A1**

- (51) International Patent Classification<sup>7</sup>: C12N 9/20 // C11D 3/386, A21D 8/04, 10/00, C11B 3/00 (74) Common Representative: NOVOZYMES A/S; Patents, Kroghshøjvej 36, DK-2880 Bagsværd (DK).
- (21) International Application Number: PCT/DK01/00600 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date:  
18 September 2001 (18.09.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
PA 2000 01411 25 September 2000 (25.09.2000) DK  
60/237,175 2 October 2000 (02.10.2000) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): NOVOZYMES A/S [DK/DK]; Kroghshøjvej 36, DK-2880 Bagsværd (DK).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): NIELSEN, Tom, Anton, Busk [DK/JP]; 186-2 Chigusacho, Hanamigawa-ku, Chiba 262-0012 (JP). ISHII, Michiyo [JP/JP]; 4-24-4, Higashi Kanamachi, Katsushika, Tokyo 125-0041 (JP). PATKAR, Shamkant, Anant [DK/DK]; Christoffers Allé 91, DK-2800 Lyngby (DK). TSUTSUMI, Noriko [JP/JP]; 3-2-16, Higashisugano, Ichikawa-shi, Chiba 272 (JP).
- Published:  
— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/24881 A1

(54) Title: PHOSPHOLIPASE FROM ZYGOASCUS HELLENICUS

(57) Abstract: The inventors discovered and isolated a phospholipase from *Zygoascus hellenicus*. The inventors also isolated the gene encoding the novel phospholipase and cloned it into an *E. coli* strain.

## PHOSPHOLIPASE FROM ZYGOASCUS HELLENICUS

## FIELD OF THE INVENTION

The present invention relates to a phospholipase, methods of using and producing it, as well as a nucleic acid sequence encoding it.

## 5 BACKGROUND OF THE INVENTION

Phospholipases are known to be useful, e.g. in baking and oil degumming.

Fu et al., Microbiology 143, 331-340 (1997) describe the cloning and characterization of a gene (LIP1) which encodes a lipase from the pathogenic yeast *Candida albicans* and state that the lipase has no phospholipase activity. The GenBank accession number for  
10 LIP1 is U34807.

LIP4 (EMBL, GenBank AF 191317) is described as a secreted lipase from *Candida albicans*.

Molecular cloning and expression of phospholipase B genes from *Candida albicans* has been described. Sugiyama et al., Medical Mycology, 37 (1), 61-67 (1999). Hoover et al.,  
15 FEMS Microbiology Letters, 167 (2), 163-169 (1998). Leidich et al., J. Biological Chemistry, 273 (40), 26078-26086 (1998). GenBank AL033501.

## SUMMARY OF THE INVENTION

The inventors discovered and isolated a phospholipase from *Zygoascus hellenicus*. The inventors also isolated and sequenced the gene encoding the novel phospholipase and  
20 cloned it into an *E. coli* strain.

Accordingly, the invention provides a phospholipase which may be a polypeptide having an amino acid sequence as the mature peptide shown in SEQ ID NO: 1 or a polypeptide encoded by the phospholipase encoding part of the DNA sequence cloned into a plasmid present in *E. coli* DSM 13648.

25 The phospholipase may also be an analogue of the polypeptide defined above which:

- i) has at least 50 % identity with said polypeptide,
- ii) is immunologically reactive with an antibody raised against said polypeptide in purified form,
- 30 iii) is an allelic variant of said polypeptide,

Finally, the phospholipase of the invention may be a polypeptide which is encoded by a nucleic acid sequence which hybridizes at 55°C with a complementary strand of the nucleic acid sequence of SEQ ID NO: 1 encoding the mature polypeptide or a subsequence thereof having at least 100 nucleotides.

The nucleic acid sequence of the invention may comprise a nucleic acid sequence which encodes the phospholipase described above, or it may encode a phospholipase and comprise:

- a) the DNA sequence encoding a mature phospholipase cloned into a plasmid present in *Escherichia coli* DSM 13648,
- b) the DNA sequence encoding a mature phospholipase shown in SEQ ID NO: 1, or
- c) an analogue of the DNA sequence defined in a) or b) which
  - i) has at least 60 % identity with said DNA sequence, or
  - ii) hybridizes at 55°C with said DNA sequence, its complementary strand or a subsequence thereof.

Other aspects of the invention provide a recombinant expression vector comprising the DNA sequence, and a cell transformed with the DNA sequence or the recombinant expression vector.

A comparison with known sequences shows that the closest prior-art sequence is LIP4 (EMBL, GenBank AF 191317), described as a secreted lipase from *Candida albicans*. The identity was calculated to be 38 % for the mature proteins and 52 % for DNA sequences encoding the mature proteins, disregarding the STOP codons.

## DETAILED DESCRIPTION OF THE INVENTION

### Genomic DNA source

A phospholipase of the invention may be derived from a strain of *Zygoascus*, particularly *Z. hellenicus*, e.g. strain CBS 4075, using probes designed on the basis of the DNA sequences in this specification. The strain is commercially available from Centraalbureau voor Schimmelcultures, Baarn, the Netherlands.

A strain of *Escherichia coli* containing a gene encoding phospholipase was deposited by the inventors under the terms of the Budapest Treaty with the DSMZ - Deutsche Sammlung von Microorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig DE, Germany. The deposit date was 10 August 2000, and the accession number was DSM 13648. The deposit was made by Novo Nordisk A/S and was later assigned to Novozymes A/S.

### Properties of phospholipase

The phospholipase has phospholipase A activity and is able to hydrolyze lecithin by releasing fatty acid. It has no lipase activity and shows no activity towards tributyrin. The isoelectric point is around 4.2.

**Recombinant expression vector**

The expression vector of the invention typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a selectable marker, a transcription terminator, a repressor gene or various activator genes.

- 5 The vector may be an autonomously replicating vector, or it may be integrated into the host cell genome.

**Production by cultivation of transformant**

- The phospholipase of the invention may be produced by transforming a suitable host cell with a DNA sequence encoding the phospholipase, cultivating the transformed  
10 organism under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

- The host organism is particularly a eukaryotic cell, in particular a fungal cell, such as a yeast cell or a filamentous fungal cell, such as a strain of *Aspergillus*, *Fusarium*, *Trichoderma* or *Saccharomyces*, particularly *A. niger*, *A. oryzae*, *F. graminearum*, *F.*  
15 *sambucinum*, *F. cerealis* or *S. cerevisiae*, e.g. a glucoamylase-producing strain of *A. niger* such as those described in US 3677902 or a mutant thereof. The production of the lysophospholipase in such host organisms may be done by the general methods described in EP 238,023 (Novo Nordisk), WO 96/00787 (Novo Nordisk) or EP 244,234 (Alko).

**Hybridization**

- 20 The hybridization is used to indicate that a given DNA sequence is analogous to a nucleotide probe corresponding to a DNA sequence of the invention. The hybridization conditions are described in detail below.

- Suitable conditions for determining hybridization between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter containing the DNA  
25 fragments or RNA to hybridize in 5 x SSC (standard saline citrate) for 10 min, and prehybridization of the filter in a solution of 5 x SSC (Sambrook et al. 1989), 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) *Anal. Biochem.* 132:6-13), <sup>32</sup>P-  
30 dCTP-labeled (specific activity > 1 x 10<sup>9</sup> cpm/µg ) probe for 12 hours at approx. 45°C. The filter is then washed two times for 30 minutes in 2 x SSC, 0.5 % SDS at a temperature of at least 55°C, more particularly at least 60°C, more particularly at least 65°C, even more particularly at least 70°C, especially at least 75°C.

- Molecules to which the oligonucleotide probe hybridizes under these conditions are  
35 detected using a x-ray film.

**Alignment and identity**

The phospholipase and the nucleotide sequence of the invention may have identities to the disclosed sequences of at least 60 % or at least 70 %, e.g. at least 85 %, particularly at least 90 % or at least 95 %, e.g. at least 98 %.

5 For purposes of the present invention, alignments of sequences and calculation of identity scores were done using the software Align, a Needleman-Wunsch alignment (i.e. global alignment), useful for both protein and DNA alignments. The default scoring matrices BLOSUM50 and the identity matrix are used for protein and DNA alignments respectively. The penalty for the first residue in a gap is -12 for proteins and -16 for DNA, while the penalty for additional residues in a gap is -2 for proteins and -4 for DNA. Align is from the  
10 FASTA package version v20u6 (W. R. Pearson and D. J. Lipman (1988), "Improved Tools for Biological Sequence Analysis", PNAS 85:2444-2448, and W. R. Pearson (1990) "Rapid and Sensitive Sequence Comparison with FASTP and FASTA", Methods in Enzymology, 183:63-98).

**15 Use of phospholipase**

The phospholipase of the invention can be used in various industrial application of phospholipases, e.g. as described below.

**Use in baking**

The phospholipase of the invention can be used in the preparation of dough and  
20 baked products (such as bread and cakes), e.g. to achieve a larger loaf volume of the baked product and/or a better shape retention during baking and/or to improve the elasticity of the baked product. Thus, the phospholipase can be used in a process for making bread, comprising adding the phospholipase to the ingredients of a dough, kneading the dough and baking the dough to make the bread. This can be done in analogy with US 4567056 or WO  
25 99/53769.

**Use in detergent**

The variant may be used as a detergent additive, e.g. at a concentration (expressed as pure enzyme protein) of 0.001-10 (e.g. 0.01-1) mg per gram of detergent or 0.001-100 (e.g. 0.01-10) mg per liter of wash liquor.

30 The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations. In a laundry detergent, the variant may be effective for the removal of fatty  
35 stains, for whiteness maintenance and for dingy cleanup. A laundry detergent composition may be formulated as described in GB 2247025, WO 9901531 or WO 9903962.

The detergent composition of the invention may particularly be formulated for hand or machine dishwashing operations. e.g. as described in GB 2,247,025 (Unilever) or WO 99/01531 (Procter & Gamble). In a dishwashing composition, the variant may be effective for removal of greasy/oily stains, for prevention of the staining /discoloration of the dishware and  
5 plastic components of the dishwasher by highly colored components and the avoidance of lime soap deposits on the dishware.

#### Other uses

The phospholipase of the invention can be used to improve the filterability of an aqueous solution or slurry of carbohydrate origin by treating it with the phospholipase. This is  
10 particularly applicable to a solution of slurry containing a starch hydrolyzate, especially a wheat starch hydrolyzate, since this tends to be difficult to filter and to give cloudy filtrates. The treatment can be done in analogy with EP 219,269 (CPC International).

Further, the phospholipase of the invention may be used for partial hydrolysis of phospholipids, particularly lecithin, to obtain improved phospholipid emulsifiers. This applica-  
15 tion is further described in Ullmann's Encyclopedia of Industrial Chemistry (Publisher: VCH Weinheim (1996)), JP patent 2794574, and JP-B 6-087751.

Further, the phospholipase of the invention may be used in a process for the production of an animal feed which comprises mixing the phospholipase with feed substances and at least one phospholipid. This can be done in analogy with EP 743 017.

20 Even further the phospholipase of the invention can be used in a process for reducing the content of phospholipid in an edible oil, comprising treating the oil with the phospholipase so as to hydrolyze a major part of the phospholipid, and separating an aqueous phase containing the hydrolyzed phospholipid from the oil. This process is applicable to the purification of any edible oil which contains phospholipid, e.g. vegetable oil such as soy bean oil,  
25 rape seed oil and sunflower oil. JP-A 2-153997, US 5264367.

The phospholipase can also be used for making cheese as described in WO 00/54601.

## **EXAMPLES**

### **Materials and methods**

#### 30 Methods

Unless otherwise stated, DNA manipulations and transformations were performed using standard methods of molecular biology as described in Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology", John Wiley and Sons,  
35 1995; Harwood, C. R., and Cutting, S. M. (eds.) .

### Enzymes

Enzymes for DNA manipulations (e.g. restriction endonucleases, ligases etc.) are obtainable from New England Biolabs, Inc. and were used according to the manufacturer's instructions.

### 5 Plasmids/vectors

pT7Blue (Invitrogen, Netherlands)

### Cloning

LA PCR™ in vitro Cloning Kit (TaKaRa) was used for cloning according to the manufacturer's instructions.

### 10 Microbial strains

*E. coli* JM109 (TOYOBO, Japan)

*E. coli* DB6507 (F, pnrF74::Tn5, seupE44, lacY1, ara-14, galK2, xyl-5, mtl-1, leuB6, proA2, hsdS20, recA13, rpsL20, thi-1, lambda-)

### 15 Media and reagents

Cove: 342.3 g/L Sucrose, 20 ml/L COVE salt solution, 10mM Acetamide, 30 g/L noble agar.

Cove-2: 30 g/L Sucrose, 20 ml/L COVE salt solution, 10mM, Acetamide, 30 g/L noble agar.

20 Cove salt solution: per liter 26 g KCl, 26 g MgSO<sub>4</sub>-7aq, 76 g KH<sub>2</sub>PO<sub>4</sub>, 50ml Cove trace metals.

Cove trace metals: per liter 0.04 g NaB<sub>4</sub>O<sub>7</sub>-10aq, 0.4 g CuSO<sub>4</sub>-5aq, 1.2 g FeSO<sub>4</sub>-7aq, 0.7 g MnSO<sub>4</sub>-aq, 0.7 g Na<sub>2</sub>MoO<sub>2</sub>-2aq, 0.7 g ZnSO<sub>4</sub>-7aq.

AMG trace metals: per liter 14.3 g ZnSO<sub>4</sub>-7aq, 2.5 g CuSO<sub>4</sub>-5aq, 0.5 g NiCl<sub>2</sub>, 13.8 g FeSO<sub>4</sub>, 8.5 g MnSO<sub>4</sub>, 3.0 g citric acid.

YPG: 4 g/L Yeast extract, 1 g/L KH<sub>2</sub>PO<sub>4</sub>, 0.5 g/L MgSO<sub>4</sub>-7aq, 5 g/L Glucose, pH 6.0.

STC: 0.8 M Sorbitol, 25 mM Tris pH 8, 25 mM CaCl<sub>2</sub>.

STPC: 40 % PEG4000 in STC buffer.

30 Cove top agarose: 342.3 g/L Sucrose, 20 ml/L COVE salt solution, 10mM Acetamide, 10 g/L low melt agarose.

MLC : per liter 40 g glucose, 50 g soybean powder, 4 g citric acid, pH 5.0.

MU-1: per liter 260 g maltodextrin, 3 g MgSO<sub>4</sub>-7aq, 6 g K<sub>2</sub>SO<sub>4</sub>, 5 g KH<sub>2</sub>PO<sub>4</sub>, 0.5 ml Trace metal solution for AMG (MU-1)

35 Trace metal solution for AMG (MU-1) : per 100 ml 1.39 g FeSO<sub>4</sub>-7aq, 1.356 g MnSO<sub>4</sub>-5aq, 0.68 g ZnCl<sub>2</sub>, 0.25 g CuSO<sub>4</sub>-5aq, 0.024 g NiCl<sub>2</sub>-6aq, 0.3 g Citric acid

MT-O medium

Soybean powder	30 g/liter
Corn steep powder	5 g/liter
Yeast extract	1 g/liter
Peptone	5 g/liter
Glucose	10 g/liter
NH <sub>4</sub> NO <sub>3</sub>	2.5 g/liter
K <sub>2</sub> HPO <sub>4</sub>	4 g/liter
MgSO <sub>4</sub> 7 H <sub>2</sub> O	0.1 g/liter
Olive oil	2 ml/shake flask
Adeka Pluronic L-61	2 drops/shake flask

Deionized water is used. pH is adjusted to 6.5 before autoclaving. 100 ml substrate/shake flask (500 ml). Adeka Pluronic L-61 is an antifoam from Asahi Denka Kogyo K.K. (Japan).

5      Lecithin plates

Substrate: 0.5 % L-alpha Phosphatidyl Choline 95% (AVANTI 441601).

Emulsifier: 0.25% Cholic acid

Buffer: 50mM acetate buffer pH5.5

10mM CaCl<sub>2</sub>

## 10      Agar Noble 2%

2% Brilliant Green 15ml/L

**Example 1: Production of phospholipase**

*Zygoascus hellenicus* CBS 4075 was grown for 4 days at 25°C on PDA slants (product of Difco with 1 % extra agar added).

15      Cells were harvested in 0.1% Tween 20 water and 57 MT-O shake flasks were inoculated and incubated on a rotary shaker at 30°C for 3 days.

Cells and particulate matter were removed by centrifugation giving 4.4 liter of supernatant. The supernatant was concentrated by ultrafiltration, washed with ca. 2 volumes of deionized water and freeze-dried.

20      About 40 g of freeze-dried powder was obtained.

**Example 2: Purification of wild type Phospholipase**

1-gram powder from Example 1 was dissolved in 100 ml water and ultra filtrated using Amicon filter (YM10 kDa cut off). After concentration to 10 ml, pH of the solution was adjusted to 8 and conductivity was 2 mSi.



The sample was then applied on anion exchanger FFQ sepharose equilibrated with 50 mM Borate buffer pH 8. Bound enzyme was eluted using linear salt gradient using same buffer containing 1 M salt. Fractions containing protein were collected.

SDS-PAGE of the fractions showed a single band.

- 5 N-terminal of the blotted protein was determined. The sequence of the N-terminal was as shown in SEQ ID NO: 8.

The enzyme was qualitatively assayed to confirm phospholipase activity using Lecithin as substrate and NEFA-kit from Waco chemicals to determine free fatty acids released.

### Example 3: Cloning and expression of lipase gene from *Zygoascus hellenicus*

#### 10 Cloning of phospholipase gene from *Zygoascus hellenicus*

A strain of *Zygoascus hellenicus* was used as a genomic DNA supplier. Cloning was carried out with adaptor PCR method using LA PCR™ in vitro Cloning Kit (TaKaRa)

- PCR reactions on *Zygoasucus hellenicus* genomic DNA which was digested by Pst I and ligated to Pst I cassette of the kit was done with ZheIN-2 (SEQ ID NO: 3) designed from  
15 N- terminal amino acid sequence of purified protein.

Reaction components (20ng / $\mu$ l of genomic DNA with cassette, 1 mM dNTP each, primer 2 nM each, 2.5mM MgCl<sub>2</sub>, 0.5 U/  $\mu$ l in LA Taq polymerase in 1X buffer (TaKaRa, Japan)) were mixed and submitted to PCR under the following conditions.

Step	Temperature	Time
1	94°C	30sec
2	50°C	2 min
3	72°C	1 min
4	4°C	forever

Steps 1 to 3 were repeated 30 times.

- 20 Amplified 640bp of the fragment was purified by GFX™ PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and sequenced.

In order to clone the missing part of the gene, genome walking was done.

- N-terminal part was obtained from the reaction mixture; 20 ng/ $\mu$ l of genomic DNA with BamH I cassette, 1 mM dNTP each, 2 nM, ZNr-2 primer, 2 nM, cassette primer, 2.5mM  
25 MgCl<sub>2</sub>, 0.5 U/  $\mu$ l in LA Taq polymerase in 1X buffer (TaKaRa, Japan).

ZNr-2: 5'-cttgtagagcagctggaagatttc-3'

Step	Temperature	Time
1	94°C	30sec
2	55°C	2 min
3	72°C	1 min
4	4°C	forever

Steps 1 to 3 were repeated 30 times.

C-terminal part was obtained by nested PCR with the primers ZC-1 and ZC-2 (SEQ ID NOS: 4 and 5) and genomic DNA with BamH I cassette. Reaction condition was same as above.

Obtained fragments were purified by GFX™ PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and sequenced with each primer which amplified the fragment.

The fidelity of LA-taq polymerase is not so good so in order to get the right sequence whole gene was amplified the primers Zhel-N (Bam) and Zhel-C (Sal) (SEQ ID NOS: 6 and 7).

Reaction components (6 ng / $\mu$ l of genomic DNA, 250 mM dNTP each, primer 250 nM each, 0.05 U /  $\mu$ l of Expand high fidelity polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for PCR under the following conditions.

Step	Temperature	Time
1	94°C	2 min
2	94°C	10sec
3	55°C	30sec
4	68°C	45sec
step 2-4 repeat 10 times		
5	94°C	10sec
6	55°C	30sec
7	68°C	45sec +20sec/cycle
step 5-7, repeat 20 times		
8	68°C	7min
7	4°C	forever

An amplified DNA fragment was gel-purified with GFX™ PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pTZhel-2, pTZhel-5, pTZhel-6, and pTZhel-7, were sequenced and all of them are identical. The sequence is defined as *Zygoascus hellenicus* phospholipase DNA sequence.

#### Expression of phospholipase gene in *Aspergillus niger*.

The phospholipase gene was digested from pTZhel-5 with BamH I and Xho I and ligated into the BamH I and Xho I sites in the *Aspergillus* expression cassette pMT2188 which has *Aspergillus niger* neutral amylase promoter, *Aspergillus nidulans* TPI leader sequences,

*Aspergillus niger* glucoamylase terminator and *Aspergillus nidulans* amdS gene as a marker and *Saccharomyces cerevisiae* URA3 gene as a marker for a plasmid construction. The ligation mixture was transformed *E. coli* 6507 by electroporation and the resultant plasmid was pZPL-8.

- 5 A strain of *Aspergillus niger* was inoculated to 100 ml of YPG medium and incubated for 16 hrs at 32°C at 120 rpm. Pellets were collected and washed with 0.6 M KCl, and resuspended in 20 ml 0.6 M KCl containing a commercial  $\beta$ -glucanase product (Glucanex, product of Novo Nordisk A/S) at the concentration of 30  $\mu$ l/ml. Cultures were incubated at 32°C at 60 rpm until protoplasts formed, then washed with STC buffer twice. The protoplasts  
10 were counted with a hemacytometer and resuspended in an 8:2:0.1 solution of STC:STPC:DMSO to a final concentration of  $2.5 \times 10^7$  protoplasts/ml. About 3  $\mu$ g of DNA was added to 100  $\mu$ l of protoplasts solution, mixed gently and incubated on ice for 30 min. One ml of SPTC was added and incubated 30 min at 37°C. After the addition of 10 ml of 50°C Cope top agarose, the reaction was poured onto Cope agar plate. Transformation  
15 plates were incubated at 32°C for 5 days.

pZPL-8 was transformed into the strain of *Aspergillus niger*. The selected transformants were inoculated in 100 ml of MLC media and cultivated at 32°C for 2 days. 5 ml of grown cell in MLC medium was inoculated to 100 ml of MU-1 medium and cultivated at 32°C for 3 days. The supernatant was obtained by centrifugation.

- 20 The phospholipase activity was checked by a lecithin plate whose composition is described above.


The substrate and the emulsifier were dissolved together in appropriate amount of D.W and autoclaved briefly (120°C for 5 min). The solution was stirred until the small particles had disappeared. Afterward it was subjected to hiscotron for 5 min. It was mixed well  
25 with other components and 25 ml aliquots were poured into Falcon 1004 plates.

Supernatant of transformants showed phospholipase activity with green halo on the assay plate.

Original (for SUBMISSION) - printed on 18.09.2001 09:06:24 AM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.92 (updated 01.03.2001)
0-2	International Application No.	PCT/DK 01/00600
0-3	Applicant's or agent's file reference	10078-WO
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	2
1-2	line	25-28
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	10 August 2000 (10.08.2000)
1-3-4	Accession Number	DSMZ 13648
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

## FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	Yes
0-4-1	Authorized officer	 Marie Louise Rosendal Head Clerk

## FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

## CLAIMS

1. A phospholipase which is:
  - a) a polypeptide encoded by the phospholipase encoding part of the DNA sequence cloned into a plasmid present in *Escherichia coli* deposit number DSM 13648,  
5 or
  - b) a polypeptide having an amino acid sequence as the mature peptide shown in SEQ ID NO: 1, or which can be obtained therefrom by substitution, deletion, and/or insertion of one or more amino acids;
  - c) an analogue of the polypeptide defined in (a) or (b) which:
    - 10 i) has at least 50 % identity with said polypeptide,
    - ii) is immunologically reactive with an antibody raised against said polypeptide in purified form,
    - iii) is an allelic variant of said polypeptide,
  - d) a polypeptide which is encoded by a nucleic acid sequence which hybridizes  
15 at 55°C with a complementary strand of the nucleic acid sequence of SEQ ID NO: 1 encoding the mature polypeptide or a subsequence thereof having at least 100 nucleotides.
2. The phospholipase of claim 1 which is native to a strain of *Zygoascus*, particularly *Z. hellenicus*, more particularly strain CBS 4075.
- 20 3. A nucleic acid sequence comprising a nucleic acid sequence which encodes the phospholipase of claim 1 or 2.
4. A nucleic acid sequence which comprises:
  - a) the partial DNA sequence encoding a mature phospholipase cloned into a plasmid present in *Escherichia coli* DSM 13648,
  - 25 b) the partial DNA sequence encoding a mature phospholipase shown in SEQ ID NO: 1,
  - c) an analogue of the sequence defined in a) or b) which encodes a phospholipase and
    - i) has at least 60 % identity with said DNA sequence, or
    - 30 ii) hybridizes at 55°C with a complementary strand of said DNA sequence or a subsequence thereof having at least 100 nucleotides,
    - iii) is an allelic variant thereof, or
  - d) a complementary strand of a), b) or c).

5. A nucleic acid construct comprising the nucleic acid sequence of claim 3 or 4 operably linked to one or more control sequences capable of directing the expression of the phospholipase in a suitable expression host.
6. A recombinant expression vector comprising the nucleic acid construct of claim 5, a promoter, and transcriptional and translational stop signals.
7. A recombinant host cell comprising the nucleic acid construct of claim 6.
8. A method for producing a phospholipase comprising cultivating the host cell of claim 7 under conditions conducive to production of the phospholipase, and recovering the phospholipase.
9. A method for preparing a dough or a baked product made from the dough, comprising adding the phospholipase of claim 1 to the dough.
10. A dough composition comprising the phospholipase of claim 1.
11. A detergent composition comprising a surfactant and the phospholipase of claim 1.
12. A process for reducing the content of phosphorus in a vegetable oil, comprising contacting the oil with the phospholipase of claim 1 in the presence of water, and then separating an aqueous phase from the oil.

10078-WO.ST25  
SEQUENCE LISTING

<110> Novozymes A/S

<120> Phospholipase

<130> 10078-WO

<160> 8

<170> PatentIn version 3.1

<210> 1

<211> 1326

<212> DNA

<213> Zygoascus hellenicus

<220>

<221> CDS

<222> (1)..(1323)

<223>

<220>

<221> mat\_peptide

<222> (52)..()

<223>

<400> 1

atg aag ttg gct gtc act ttt ctc gtt agc ttg ata tct cta gcc tgg	48
Met Lys Leu Ala Val Thr Phe Leu Val Ser Leu Ile Ser Leu Ala Trp	
-15 -10 -5	

gca ggt att cct gta gct atc tct gat gat cct tgg tat aat gct cca	96
Ala Gly Ile Pro Val Ala Ile Ser Asp Asp Pro Trp Tyr Asn Ala Pro	
-1 1 5 10 15	

gct aat ctt gat tca ctc gct gat ggt act att atc aga tct cga gaa	144
Ala Asn Leu Asp Ser Leu Ala Asp Gly Thr Ile Ile Arg Ser Arg Glu	
20 25 30	

aca ccc aac tct ctt gct gca ttg gtg ctg ttc cct caa aac gtc cag	192
---	-----

10078-WO.ST25															
Thr	Pro	Asn	Ser	Leu	Ala	Ala	Leu	Val	Leu	Phe	Pro	Gln	Asn	Val	Gln
		35						40					45		
aaa	tct	tac	cag	ctg	ctc	tac	aag	tct	act	gat	gat	ctg	aat	aga	act
Lys	Ser	Tyr	Gln	Leu	Leu	Tyr	Lys	Ser	Thr	Asp	Asp	Leu	Asn	Arg	Thr
		50					55					60			
240															
ttc	gct	aca	gca	acc	acg	atc	att	ggt	ccc	tat	aac	gct	gat	tat	acc
Phe	Ala	Thr	Ala	Thr	Thr	Ile	Ile	Val	Pro	Tyr	Asn	Ala	Asp	Tyr	Thr
	65					70					75				
288															
aag	ctt	gtc	tcc	ttc	cag	att	gca	gag	gat	tcg	gcc	tat	atc	aac	tgt
Lys	Leu	Val	Ser	Phe	Gln	Ile	Ala	Glu	Asp	Ser	Ala	Tyr	Ile	Asn	Cys
	80				85					90					95
336															
gca	ccc	tct	tat	gcc	ttg	caa	ttt	gcc	tca	gac	cct	gat	gct	ctt	ctt
Ala	Pro	Ser	Tyr	Ala	Leu	Gln	Phe	Ala	Ser	Asp	Pro	Asp	Ala	Leu	Leu
				100					105					110	
384															
tca	cag	gtt	gaa	ttt	ggt	ttc	cta	tct	gct	atg	ctt	tca	ctg	gga	tat
Ser	Gln	Val	Glu	Phe	Val	Phe	Leu	Ser	Ala	Met	Leu	Ser	Leu	Gly	Tyr
			115					120					125		
432															
gtt	gtg	aat	gtt	cct	gat	tac	gaa	ggt	ccg	cga	tct	gct	ttc	act	gta
Val	Val	Asn	Val	Pro	Asp	Tyr	Glu	Gly	Pro	Arg	Ser	Ala	Phe	Thr	Val
		130					135					140			
480															
gga	cag	gtt	gcc	ggc	aga	gca	gtc	ttg	gac	tct	ctt	aga	gca	gct	ctt
Gly	Gln	Val	Ala	Gly	Arg	Ala	Val	Leu	Asp	Ser	Leu	Arg	Ala	Ala	Leu
	145					150					155				
528															
gca	tcc	tcg	tct	att	aca	gga	atc	aat	tct	aaa	gct	cgt	att	gcc	ttg
Ala	Ser	Ser	Ser	Ile	Thr	Gly	Ile	Asn	Ser	Lys	Ala	Arg	Ile	Ala	Leu
	160				165					170					175
576															
tgg	ggc	tac	tct	ggt	ggc	tcc	ctt	gct	agt	ggc	tgg	gca	gct	cag	ctt
Trp	Gly	Tyr	Ser	Gly	Gly	Ser	Leu	Ala	Ser	Gly	Trp	Ala	Ala	Gln	Leu
				180					185					190	
624															
caa	ccc	tcg	cat	gcc	cct	gag	ttg	gtt	atc	tct	ggt	gct	gct	ttg	ggt
Gln	Pro	Ser	His	Ala	Pro	Glu	Leu	Val	Ile	Ser	Gly	Ala	Ala	Leu	Gly
			195					200					205		
672															
ggt	gtt	gta	cct	aat	att	gct	gct	gtc	gca	acc	aag	gtc	agc	ggt	ggt
Gly	Val	Val	Pro	Asn	Ile	Ala	Ala	Val	Ala	Thr	Lys	Val	Ser	Gly	Gly
		210					215					220			
720															
gta	ttc	gct	ggc	ttg	gtt	cct	gca	ggt	atc	acc	ggt	ttt	ggt	aac	cag
Val	Phe	Ala	Gly	Leu	Val	Pro	Ala	Gly	Ile	Thr	Gly	Phe	Gly	Asn	Gln
		225				230					235				
768															
tat	gat	gaa	gtt	agt	gaa	atc	att	tct	gct	cac	ttg	gtt	cct	tcc	aag
Tyr	Asp	Glu	Val	Ser	Glu	Ile	Ile	Ser	Ala	His	Leu	Val	Pro	Ser	Lys
	240				245					250					255
816															
gtt	gaa	gag	tgg	aac	aaa	gct	aat	agt	cag	tgt	atc	gtt	gct	gat	gct
Val	Glu	Glu	Trp	Lys	Ala	Asn	Ser	Ser	Gln	Cys	Ile	Val	Ala	Asp	Ala
				260					265					270	
864															
tta	gga	tat	gca	ttc	cag	gat	att	ttc	gag	gat	tat	ttc	act	att	ggt
Leu	Gly	Tyr	Ala	Phe	Gln	Asp	Ile	Phe	Glu	Asp	Tyr	Phe	Thr	Ile	Gly
			275					280					285		
912															
act	ggt	att	ctc	gag	gtt	cct	gaa	gtt	gcc	gca	aga	ctc	aat	caa	aac
Thr	Gly	Ile	Leu	Glu	Val	Pro	Glu	Val	Ala	Ala	Arg	Leu	Asn	Gln	Asn
		290					295					300			
960															
ctt	atg	ggt	tct	ttg	gtt	cct	gaa	ttt	cct	att	tac	att	tac	cat	gct
1008															



10078-WO.ST25

Leu	Met	Gly	Ser	Leu	Val	Pro	Glu	Phe	Pro	Ile	Tyr	Ile	Tyr	His	Ala	
305						310					315					
gtt	ggc	gac	gaa	gtt	gct	cct	gtc	gaa	aaa	gct	gat	act	ctt	gtc	tct	1056
Val	Gly	Asp	Glu	Val	Ala	Pro	Val	Glu	Lys	Ala	Asp	Thr	Leu	Val	Ser	
320					325					330					335	
tcc	tgg	tgc	aat	gct	ggg	atc	aaa	tct	ctt	gaa	tat	cat	aag	gat	att	1104
Ser	Trp	Cys	Asn	Ala	Gly	Ile	Lys	Ser	Leu	Glu	Tyr	His	Lys	Asp	Ile	
				340					345					350		
ttg	gat	gag	cat	ctt	tca	gag	gag	gtg	ttt	ggg	tca	gct	ggc	gcc	att	1152
Leu	Asp	Glu	His	Leu	Ser	Glu	Glu	Val	Phe	Gly	Ser	Ala	Gly	Ala	Ile	
			355					360					365			
gtc	tgg	ttg	aaa	tcc	atc	atg	gaa	gac	tct	gct	gat	ctc	gct	gat	ggc	1200
Val	Trp	Leu	Lys	Ser	Ile	Met	Glu	Asp	Ser	Ala	Asp	Leu	Ala	Asp	Gly	
		370					375					380				
tgc	aaa	act	gaa	atc	tct	ctt	tcg	caa	ctt	att	gat	cct	gat	gct	gtt	1248
Cys	Lys	Thr	Glu	Ile	Ser	Leu	Ser	Gln	Leu	Ile	Asp	Pro	Asp	Ala	Val	
	385					390					395					
cct	gtt	ctt	ggg	gaa	gag	ttg	gtg	caa	gtg	ttg	ctt	aac	tta	gtt	gga	1296
Pro	Val	Leu	Gly	Glu	Glu	Leu	Val	Gln	Val	Leu	Leu	Asn	Leu	Val	Gly	
400				405					410						415	
aag	cct	atc	ggg	ccc	atc	tcg	ttt	gcc	taa							1326
Lys	Pro	Ile	Gly	Pro	Ile	Ser	Phe	Ala								
				420												

&lt;210&gt; 2

&lt;211&gt; 441

&lt;212&gt; PRT

&lt;213&gt; Zygoascus hellenicus

&lt;400&gt; 2

Met Lys Leu Ala Val Thr Phe Leu Val Ser Leu Ile Ser Leu Ala Trp  
 -15 -10 -5

Ala Gly Ile Pro Val Ala Ile Ser Asp Asp Pro Trp Tyr Asn Ala Pro  
 -1 1 5 10 15

Ala Asn Leu Asp Ser Leu Ala Asp Gly Thr Ile Ile Arg Ser Arg Glu  
 20 25 30

Thr Pro Asn Ser Leu Ala Ala Leu Val Leu Phe Pro Gln Asn Val Gln  
 35 40 45

Lys Ser Tyr Gln Leu Leu Tyr Lys Ser Thr Asp Asp Leu Asn Arg Thr  
 50 55 60

Phe Ala Thr Ala Thr Thr Ile Ile Val Pro Tyr Asn Ala Asp Tyr Thr  
 65 70 75

10078-WO.ST25

Lys Leu Val Ser Phe Gln Ile Ala Glu Asp Ser Ala Tyr Ile Asn Cys  
80 85 90 95

Ala Pro Ser Tyr Ala Leu Gln Phe Ala Ser Asp Pro Asp Ala Leu Leu  
100 105 110

Ser Gln Val Glu Phe Val Phe Leu Ser Ala Met Leu Ser Leu Gly Tyr  
115 120 125

Val Val Asn Val Pro Asp Tyr Glu Gly Pro Arg Ser Ala Phe Thr Val  
130 135 140

Gly Gln Val Ala Gly Arg Ala Val Leu Asp Ser Leu Arg Ala Ala Leu  
145 150 155

Ala Ser Ser Ser Ile Thr Gly Ile Asn Ser Lys Ala Arg Ile Ala Leu  
160 165 170 175

Trp Gly Tyr Ser Gly Gly Ser Leu Ala Ser Gly Trp Ala Ala Gln Leu  
180 185 190

Gln Pro Ser His Ala Pro Glu Leu Val Ile Ser Gly Ala Ala Leu Gly  
195 200 205

Gly Val Val Pro Asn Ile Ala Ala Val Ala Thr Lys Val Ser Gly Gly  
210 215 220

Val Phe Ala Gly Leu Val Pro Ala Gly Ile Thr Gly Phe Gly Asn Gln  
225 230 235

Tyr Asp Glu Val Ser Glu Ile Ile Ser Ala His Leu Val Pro Ser Lys  
240 245 250 255

Val Glu Glu Trp Asn Lys Ala Asn Ser Gln Cys Ile Val Ala Asp Ala  
260 265 270

Leu Gly Tyr Ala Phe Gln Asp Ile Phe Glu Asp Tyr Phe Thr Ile Gly  
275 280 285

Thr Gly Ile Leu Glu Val Pro Glu Val Ala Ala Arg Leu Asn Gln Asn  
290 295 300

Leu Met Gly Ser Leu Val Pro Glu Phe Pro Ile Tyr Ile Tyr His Ala  
305 310 315

Val Gly Asp Glu Val Ala Pro Val Glu Lys Ala Asp Thr Leu Val Ser  
320 325 330 335

Ser Trp Cys Asn Ala Gly Ile Lys Ser Leu Glu Tyr His Lys Asp Ile  
340 345 350

10078-WO.ST25  
 Leu Asp Glu His Leu Ser Glu Glu Val Phe Gly Ser Ala Gly Ala Ile  
                   355                  360                  365

Val Trp Leu Lys Ser Ile Met Glu Asp Ser Ala Asp Leu Ala Asp Gly  
                   370                  375                  380

Cys Lys Thr Glu Ile Ser Leu Ser Gln Leu Ile Asp Pro Asp Ala Val  
                   385                  390                  395

Pro Val Leu Gly Glu Glu Leu Val Gln Val Leu Leu Asn Leu Val Gly  
                   400                  405                  410                  415

Lys Pro Ile Gly Pro Ile Ser Phe Ala  
                                   420

<210> 3

<211> 23

<212> DNA

<213> Artificial

<220>

<221> misc\_feature

<223> ZheIn-2

<220>

<221> misc\_feature

<222> (9)..(21)

<223> inosine

<220>

<221> modified\_base

<222> (9)..(9)

<223> I

<220>

<221> modified\_base

<222> (21)..(21)

<223> I

## 10078-WO.ST25

<400> 3  
gaygayccnt ggtayaaygc ncc

23

<210> 4

<211> 24

<212> DNA

<213> Artificial

<220>

<221> misc\_feature

<223> ZC-1

<400> 4  
tattgccttg tggggctact ctgg

24

<210> 5

<211> 24

<212> DNA

<213> Artificial

<220>

<221> misc\_feature

<223> ZC-2

<400> 5  
tattgccttg tggggctact ctgg

24

<210> 6

<211> 36

<212> DNA

<213> Artificial

<220>

<221> misc\_feature

<223> ZheI-N (Bam)

<400> 6  
cgcggatcca tgaagttggc tgtcactttt ctcggt

36

10078-WO.ST25

&lt;210&gt; 7

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; zhe1-c (sal)

&lt;400&gt; 7

acgcgtcgac ttaggcaaac gagatgggcc cgatagg

37

&lt;210&gt; 8

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Zygoascus hellenicus

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; N-terminal

&lt;400&gt; 8

Gly Ile Pro Met Ala Ile Ser Asp Asp Pro Trp Tyr Asn Ala Pro Ala  
1 5 10 15

Asn Leu Asp Ser Leu Ala  
20

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 01/00600

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/20 //C11D3/386,A21D8/04,A21D10/00,C11B3/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1991:118793, vol. 28, no. 5, pages 355-362, MAGO N ET AL : "Subcellular localization of enzymes of phospholipid metabolism in candida-albicans" XP002902240 abstract & JOURNAL OF MEDICAL AND VETERINARY MYCROLOGY, 1990,	1-12
A	WO 99 03962 A (BAECK ANDRE CESAR ) 28 January 1999 (1999-01-28) page 3 -page 5 --- -/--	11



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

18 January 2002

Date of mailing of the international search report

22 02 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Yvonne Siösteen

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 01/00600

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 26057 A (NOVONORDISK AS) 18 June 1998 (1998-06-18) page 14 -page 21 page 45 -page 49; claims 65-68 -----	9,10,12

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
**PCT/DK 01/00600**

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9903962	A	28-01-1999	WO 9903962 A1	28-01-1999
			AU 3884597 A	10-02-1999
			BR 9714808 A	11-12-2001
			EP 1005520 A1	07-06-2000
			JP 2001510233 T	31-07-2001
<hr/>				
WO 9826057	A	18-06-1998	AU 5187898 A	03-07-1998
			CN 1245532 A	23-02-2000
			WO 9826057 A1	18-06-1998
			EP 0869167 A2	07-10-1998
			JP 2000507458 T	20-06-2000
			US 6143545 A	07-11-2000
			US 6103505 A	15-08-2000
			EP 0884524 A2	16-12-1998
<hr/>				